



ISSN: 1813-1638

The Medical Journal of Tikrit University

Available online at: www.mjotu.com

العراقية
المجلات الاكاديمية العلمية
IRAQI
Academic Scientific Journals

Zeyad M. Sadeq ⁽¹⁾
Nadine A. Mahmood ⁽²⁾

(1) Department of
pediatrics, Medical college,
Hawler Medical University.
Iraq

(2) Department of
pediatrics, Medical college,
Hawler Medical University.
Iraq

Keywords:

*congenital heart disease,
pulmonary stenosis,
valvuloplasty*

ARTICLE INFO

Article history:

Received 05 Sep 2018
Accepted 01 Oct 2018
Available online 01 Dec 2021

**Long-term results of balloon pulmonary valvuloplasty
in pediatric age group in surgical specialty teaching
hospital/cardiac center/Hawler**

ABSTRACT

Congenital valvular pulmonary stenosis accounts for 5–10% of all congenital heart disease. Balloon pulmonary valvuloplasty is a safe and effective method in reduction of moderate to severe valvular pulmonary stenosis. This retrospective study done in Hawler surgical specialty cardiac center on pediatric patients who had pulmonary valve stenosis (with echocardiographic peak gradient ≥ 50 mmHg) and treated by balloon pulmonary valvuloplasty.

A total of 178 cases were involved, 93(52.25%) patient was male and 85 (47.75%) patients was female. 160 patient (89.89 %) were treated with single balloon method 18 patient (10.11 %) were managed by double balloon method technique. Follow up period of these patients range from 1 year to 10 years (mean of 5.5 year). In 153 patients (85.96%) pressure across pulmonary valve were below 36 mm Hg, while in 21 (11.80%) pressure gradient across pulmonary valve were (36-50 mm Hg). Four (2.24%) patients had pressure gradient across pulmonary valve more than 50 mm Hg. Forty patients (22.47%) developed mild pulmonary valve regurgitation, while 20 patient (11.24%) developed significant pulmonary valve regurgitation.

Balloon pulmonary valvuloplasty provides long-term relief of pulmonary valvular obstruction in the majority of pediatric patients. The long-term results are favorable.

DOI: <http://dx.doi.org/10.25130/mjotu.27.2021.04>

*Corresponding author E mail : ZeyadMS89@gmail.com

Introduction:

Congenital cardiac abnormality occurs in approximately 0.8% of live births. Congenital heart defects remains the leading cause of death in children with congenital malformations.¹

Obstruction to the outflow from the right ventricle, whether within the body of the right ventricle, at the pulmonary valve, or in the pulmonary arteries, is described as pulmonary stenosis.²

Pulmonary stenosis is a common congenital cardiac lesion. The incidence of valvar stenosis has been reported at 0.6 to 0.8 per 1000 live births. Balloon valvuloplasty is the procedure of choice for the valvular stenosis at all ages.³

AIM OF THE STUDY:

Is to assess the long-term results of percutaneous balloon valvuloplasty at surgical specialty hospital/ cardiac center/ Hawler).

METHODS:

This retrospective, single center, observational study involve pediatric patients (whom age range between one day up to 21 years) with history of pulmonary valve stenosis that treated

by transcatheter pulmonary valve dilatation in surgical specialty teaching hospital/cardiac center/Hawler from 1st January 2007 to 31th December 2016.

The criterion for indicating percutaneous valvuloplasty was a peak pressure gradient 50 mmHg across the pulmonary valve with normal cardiac function and right ventricular/left ventricular pressure ratio > 0.8.

Exclusion criteria include patients whom been treated with surgical dilatation, presence of pulmonary stenosis at other levels (infundibular stenosis, supra-ventricular stenosis), and pulmonary stenosis in association with other congenital heart disease.

Echo-Doppler examination using (VIVED E9) generation machine were performed on follow up. A 3MHz and 5 MHz probe were used for examination. A maximum peak instantaneous pressure gradient across RVOT was determined using the modified Bernoulli equation. $\Delta P = 4(V_2^2 - V_1^2)$

Immediate success is defined as a pulmonary gradient < 36 mmHg

following the procedure, the effectiveness of the procedure was defined by immediate success and by a subsequent decrease in transvalvular gradient to < 36 mmHg. Restenosis was defined as an increase in pressure gradient increasing to ≥ 36 mmHg after a successful procedure.⁴

Patients' medical records were reviewed and the following procedural data were collected: age, weight at the time of valvuloplasty; clinical presentation; valve morphology, pre- and post valvuloplasty hemodynamic data (pulmonary transvalvular gradient, RV pressure, and right ventricular/left ventricular pressure ratio. Pulmonary regurgitation assessed by the ratio between the regurgitant jet and pulmonary annulus diameters, establishing 4 grades (grade I, ratio 10%; grade II, 11% to 25% grade III, 26% to 50% and grade IV $> 50\%$)².

Finally, the length of follow-up was analyzed, defined as the time from valvuloplasty until final echocardiographic follow-up

Chi square Test were used for analyses.

P value < 0.05 was considered significant for all tests.

RESULTS:

A total of 178 cases were involved in the study; 93(52.25%) patient was male and 85 (47.75%) patients was female, male to female ratio was 1.1/1.

Age of the patients range from 17 days old to 21 years, with mean age of (4.06 ± 4.7) years old. Pressure gradient across the pulmonary valve range from 50mm Hg to 179mm Hg with mean pressure of (98.98 ± 24.63) . Follow up period of these patients range from 1 year to 10 years (mean of 5.5 year), (table1).

The morphology of the pulmonary valve prior to transcatheter intervention was varied between domed valve found in 127(71.35%) case, dysplastic valve in 42 (23.60) cases, while the remaining 9 (5.05%) cases were have more complex valve morphology (table 2).

Regarding treatment modality, it was recorded that 18 patient (10.11 %) were managed by double balloon method technique, the remaining 160 patient (89.89 %) were treated with

single balloon method, (table 3). Studied group were divided into two subgroups according to the early result of the balloon valvuloplasty, In 155 patient (87.08 %); the immediate post valvuloplasty decrement in the pressure gradient across pulmonary valve were below 36 mm Hg and these patient had successful balloon dilatation(group one). In 23 patient (12.92 %); balloon dilatation were only partially successful (pressure gradient across the pulmonary valve were decrease between 36-50 mm Hg)(group two) as shown in table (4). Regarding the results of long term follow up, it was found that among the patients of group one, (137 patients 88.39 %) still has transvalvular pressure gradient less than 36 mm Hg across pulmonary valve, while the pressure gradient remain more than 36 mm Hg in 18 patients (11.61 %). regarding

patients in group two, sixteen patients (65.22 %), show decline in the pressure gradient across the pulmonary valve (transvalvular pressure less than 36 mm Hg), while pressure gradient remain between (36-50 mm Hg) in three patients (13.04 %). In the remaining 4 patients (21.74 %) severe stenosis that demand surgical intervention to relieve transvalvular obstruction.

Grade 4 pulmonary valve regurgitation was found 17 (10.96 %) patients of group one, while 53 (22.58 %) patients found to have grade 2-3 pulmonary regurgitation. Regarding group two; 3 (12.5%) developed significant (grade 4) regurgitation, while Grade 2-3 pulmonary valve regurgitation found in 5 (20.8 %) patient in group two.

Table (1): Characters of patients whom involve in the study

Parameter		
Age	0-21 years	4.06 ±4.7
Pressure gradient	50-179 mm Hg	98.98±24.63
Period of follow up	1-10 years	4.5±2.7

Females	85 (47.75 %)	
Males	93 (52.25 %)	

Table (2): morphology of the pulmonary valve prior to transcatheter intervention

Morphology	No.	%
Domed	127	71.35
Dysplastic	42	23.60
Complex	9	5.05
Total	178	100

Table (3): Methods of balloon dilatation used in pulmonary valvuloplasty

	No. of patients	%
Single balloon	160	89.89
Double balloon	18	10.11
Total	178	100

Table (4): immediate results of post transcatheter pulmonary valve dilatation.

Results of intervention	No.	%
Successful (group one)	155	87.08
Partially successful (group two)	23	12.92
Total	178	100

Table (5): Long term results of balloon pulmonary valvuloplasty-pressure gradient across pulmonary valve

Transvalvular pulmonary pressure	Group one	Group two	Total
Less than 36 mm Hg	137 (88.39 %)	16 (69.56 %)	153 (85.96%)

36-50 mm Hg	18 (11.61 %)	3 (13.04 %)	21 (11.80%)
More than 50 mm Hg	--	4 (17.39 %)	4 (2.24%)
	155 (100 %)	23 (100 %)	178 (100%)

P value < 0.05

Table (6): Long term results of balloon pulmonary valvuloplasty- pulmonary valve regurgitation

	Group one	Group two	Total
Mild (Grade 2-3) regurgitation	35 (22.58 %)	5 (20.8 %)	40 (22.47%)
Significant (grade 4) regurgitation	17 (10.96 %)	3 (12.5 %)	20 (11.24%)
Total	52 (33.54 %)	8 (32.33 %)	60 (33.71%)

P value > 0.05

DISCUSSION:

Percutaneous balloon pulmonary valvuloplasty (BPV) represents a major milestone in the field of interventional cardiology, regardless of the patient's age and valve morphology; BPV has replaced surgery as the initial treatment of choice for patients with moderate to severe pulmonary valve stenosis.⁵

In this study, 18 patient (10.11 %) were managed by double balloon method

technique, the remaining 160 patient (89.89 %) were treated with single balloon method. In a similar study that included 118 patients with congenital valvular pulmonary stenosis at Jipmer tertiary care hospital in India from 1988 to 2011, Ezhumalai, et al found that double balloon technique was used in only 1 patient (0.9%)⁶. Simultaneous double balloon dilation has the advantage of smaller profiles that

minimize sheath requirements and potential vascular injury, although they require multiple sites of vascular entry. There are limited data evaluating the unequal distribution of wall stresses that may occur with this technique although no untoward clinical effects have been reported.⁷

Results of double balloon valvuloplasty are excellent but not superior to single balloon valvuloplasty, double balloon valvuloplasty should be used if the pulmonary valve annulus is too large to dilate with a single balloon. There is no need for double balloon valvuloplasty if the pulmonary valve can be dilated with a commercially available single balloon.⁸

Regarding long term outcome of the percutaneous pulmonary valve dilatation Overall successfulness in this study was 153 (85.96%), these patients shows pressure gradient across pulmonary valve below 36 mm Hg taken by doppler echocardiography, and in 21 (11.80%) patient found to have pressure gradient between (36-50 mm Hg). also its found that immediate post

dilatation results have a great influence on the expected long term outcome since statistically significant difference in the rate of success exist between those who shows full successfulness and those who shows only partial successfulness (88.39 % Vs 69.56 %). In a study done at The Johns Hopkins Hospital in Between June 1981 and December 1986, initial BPV was attempted in 49 patients. 44 patients in whom immediate result fall below 36 mmHg Only, 2 patients (4.55 %) shows pressure gradient over 36 mm Hg. the remaining 42 patients (94.45 %) shows pressure gradient below 36 mmHg. in another 5 patient (10.2%) involved in the mentioned study; pressure gradient after immediate post catheterization period fail to decline below 36 mm Hg; long term study of this group shows that one patient (20%) undergone surgical dilatation, one patient (20%) did re catheterization dilatation, and in 3 (60%) patients pressure gradient decline below 36 mm Hg. ⁹. In Ezhumalai, et al study ⁶ the long term success rate (≥ 6 years after

valvuloplasty) was 99(91.7%), and 2(1.9 %) patient developed significant restenosis. Raquel Merino study 53 patients diagnosed with pulmonary valve stenosis who underwent percutaneous balloon pulmonary valvuloplasty between December 1985 and December 2000 in Virgen del Rocío tertiary care Hospital, Seville, Spain center, occurrence of long-term restenosis in 1 patient (2.1%)¹⁰.

Although the result of all these studies appeared to be encouraging, differences in the success rate may partially related to time of presentation of the patient, availability of catheterization equipment (including proper balloon size, and regular follow up regimen).

Regarding pulmonary insufficiency, 40 patients (22.35 %) have grade 2-3 pulmonary, and in 20 patients (11.17 %) echocardiographic study shows grade (grade 4) insufficiency. No statistical differences found in the degree of pulmonary regurgitation between patients that shows full immediate successfulness and those with partial successfulness after balloon

dilatation. A study done by Garty Y et al. in the Cardiac Catheterization Laboratories, Department of Cardiology, Hospital for Sick Children, in Toronto Canada, 150 children who had PBD between 1984 and 1992 were reviewed. Moderate or severe PR found in 85 patient (57%) of children on long term follow-up.¹¹

In a similar study done in kingdom of Saudi showed that 26 of 35 (74%) patients with successful balloon pulmonary valvuloplasty developed pulmonary insufficiency on long-term follow-up. This is understandable, because the balloon valvuloplasty produces commissural splitting and tearing or avulsion of valve leaflets.¹²

In a study done in Tunisia, Fattouma Bourguiba University Hospital. From April 1987 to April 1996, 66 patients underwent percutaneous BPV in catheterization laboratory for relief of isolated pulmonary valve stenosis. Mild to moderate pulmonary regurgitation was present in 39% of patients.¹³

In a study done in Iran Shahid Sadoughi University of Medical Sciences,. From

June 1998 to January 2012 sixty consecutive patients (33 females, 27 males) with moderate to severe valvar pulmonary stenosis. Pulmonary insufficiency was demonstrated in 17 (32%) at late follow up. A second valvuloplasty was performed in two (3.8%) patients presenting with re-stenosis.¹⁴

Although the result of all studies are good enough to consider balloon pulmonary valvuloplasty as a gold standard of treatment in congenital pulmonary stenosis, the deference in the long term follow up probably due to the slandered of the centers in which the catheterization carried in, together with the accumulative experience of the physician that did the catheterization, since the studies are done over deferent time.

CONCLUSION:

The long-term results of balloon valvuloplasty for pulmonary stenosis appear encouraging. Repeat catheterization and echo doppler data suggest persistent relief of pulmonary

stenosis. Less than moderate pulmonary regurgitation develops in most cases, but rarely requires surgical treatment.

REFERENCES:

1. Bernstein D. Epidemiology and Genetic Basis of Congenital Heart Disease. In: Kliegman RM, Behrman RE, Jenson HB Stanton BF. Nelson Textbook of Pediatrics. 18th ed. Philadelphia: WB Saunders, 2007:1499
2. Schneider DS. Acyanotic Congenital Heart Disease. In: Kliegman, RM, Karen JM, Maccdante KJ, Jenson HB. Nelson Essentials of Pediatrics, 6th ed. Canada: WB Saunders, 2011: 537-46.
3. Keane JF, Fyler DC. Pulmonary Stenosis Alexander ME, Berul CI, Blume ED , Breitbart RE, Cecchin F, Clapham DE et al Nadas' Pediatric Cardiology. 2nd ed Philadelphia: WB Saunders, 2006:549-56
4. Jarrar M, Betbout F, Farhat MB, Maatouk F, Gamra H, Addad F, et al. Long-term invasive and noninvasive results of percutaneous balloon pulmonary valvuloplasty in children, adolescents, and adults. Am Heart J. 1999;138(5 Pt 1):950-4.
5. Mullins CE, Latson LA, Neches WH,

- Colvin EV, Kan J: Balloon dilation of miscellaneous lesions: Results of Valvuloplasty and Angioplasty of Congenital Anomalies Registry. *Am J Cardiol* 1990;65:802-803.
6. Ezhumalai B, Awasthy N, Ananthakrishna A, Satheesh S, Jayaraman B (2016) Long-Term Follow-Up Analysis of Percutaneous Balloon Pulmonary Valvuloplasty: Experience of an Indian Tertiary Care Hospital. *J Clin Trial Cardiol* 3(1): 1-5.
7. Al Kasab S, Ribeiro P, Al Zaibag M. Use of double balloon technique for percutaneous balloon pulmonary valvotomy in adults. *Br Heart J*. 1987;58(2):136-141.
8. Rao Psyamasundar, Fawzy Mohammed. Double Balloon Technique for Percutaneous Balloon Pulmonary Valvuloplasty: Comparison with Single Balloon Technique. *Journal of Interventional Cardiology*. 1988;1(4):257-262
9. Brian W. McCrindle, MD, MPH, and Jean S. Kan, MD Long-term Results After Balloon Pulmonary Valvuloplasty. *Invasive Cardiol*. 2005 Jun;17(6):323-5.
11. Raquel Merino-Ingelmo, Jose' Santos-de Soto, Fe'lix Coserria-Sa' nchez, Alfonso Descalzo-Sen'oran, and Israel Valverde-Pe'rez: Long-term Results of Percutaneous Balloon Valvuloplasty in Pulmonary Valve Stenosis in the Pediatric Population. *Rev Esp Cardiol*. 2014;67(5):374-379.
12. Walls JT, Lababidi Z, Curtis JJ, Silver D. Assessment of percutaneous balloon pulmonary and aortic valvuloplasty. *J Thorac Cardiovasc Surg* 1984;88:352-6.
13. Mourad J, Fethi B, Mohamed BF. Long-term invasive and noninvasive results of percutaneous balloon pulmonary valvuloplasty in children, adolescents and adult. *American Heart Journal* 1999 Dec; 138(5 Pt 1):950-4.
14. Behjati AM, Moshtaghion SH, Rajaei S; Long-term Results of Balloon Pulmonary Valvuloplasty in Children with Congenital Pulmonary Valve Stenosis, *Iranian Journal of Pediatrics* 2013 Feb; 23 (1) 32-36.
10. [Garty Y](#), [Veldtman G](#), [Lee K](#), [Benson L](#) : Late outcomes after pulmonary valve balloon dilatation in neonates, infants and children. *J*